

The investigation of the $\gamma\gamma \rightarrow W^+W^-$ including electromagnetic corrections at the TESLA kinematics

Author I.B. Marfin

NCHEP

153 Bogdanovitcha str., 220040 Minsk, Belarus

E-mail: marfin@hep.by

Author V.A. Mossolov

NCHEP

153 Bogdanovitcha str., 220040 Minsk, Belarus

E-mail: mos@hep.by

Author T.V. Shishkina

NCHEP

153 Bogdanovitcha str., 220040 Minsk, Belarus

E-mail: shishkina@hep.by

Abstract.

The W^+W^- production in $\gamma\gamma$ scattering is considered in the Standard Model. The main contribution to radiative effects for the process $\gamma\gamma \rightarrow W^+W^-$ has been calculating and analyzing. It is found the latter is considerable at high energies and greatly contributes to the differential cross section $d\sigma_{\lambda_1\lambda_2\lambda_3\lambda_4}$ at various polarizations of initial photons and final bosons. Monte-Carlo generator built based on TESLA kinematics.

1. Introduction

The multiple vector-boson production will be a crucial test of gauge structure of the Standard Model since the triple and quartic vector-boson couplings involved in this kind of reaction are constrained by the $SU(2) \otimes U(1)$ gauge invariance. The production of several vector bosons is the best place to search directly for any anomalous behavior of triple and quartic couplings [1].

Any small deviation from the Standard Model predictions for these coupling spoils the cancellations of the high energy behaviour between the various diagrams, giving rise to an anomalous growth of the cross section with energy. It is important to measure the vector-boson selfcouplings and look for deviations from the Standard Model.

Using the linear e^+e^- -collider, one can obtain the colliding γe and $\gamma\gamma$ beams with the same energies as in e^+e^- collisions and with high luminosity.

Thus, there is the possibility of direct experimental investigation of the reactions:

$$\gamma e \rightarrow W\nu$$

and

$$\gamma\gamma \rightarrow W^+W^-.$$

These reactions were considered as parts of the W-bosons production process in e^+e^- collisions ($ee \rightarrow eW\nu, e^+e^- \rightarrow W+X, ee \rightarrow eeWW$) and in ep and pp collisions as well. However, the corresponding cross sections are very small and such processes will be observed in the nearest future hardly.

Reactions $e^+e^- \rightarrow W^+W^-$ and $\gamma\gamma \rightarrow W^+W^-$ will give the unique possibilities of W -investigation due to relatively large cross sections and low background. Moreover, the colliding γe and $\gamma\gamma$ beams will allow us to study a number of problems which are harding solved in other collisions [2]. The cross sections of process $\gamma\gamma \rightarrow W^+W^-$ is of order $\sim 4\pi\alpha^2/M_W^2$ at high energy (they are constant at $s \rightarrow \infty$ due to the t-channel vector exchange). Therefore, at $s_{\gamma e} \sim s_{\gamma\gamma} \sim s_{ee} \gg M_W^2$, the cross sections of this reaction considerably exceed the cross section of the e^+e^- annihilation into hadrons.

2. The probing triple and quartic gauge bosons couplings

The cross section $\gamma\gamma \rightarrow W^+W^-$ is about an order of magnitude larger than $e^+e^- \rightarrow W^+W^-$.

We discuss how one can use $\gamma\gamma$ collisions to study the pair production of vector bosons (see fig. 1)

$$\gamma + \gamma \rightarrow W^+ + W^-$$

and get information on three- and four-boson vertices.

The process $\gamma\gamma \rightarrow W^+W^-$ depends at the tree level both on the triple γWW and the quartic $\gamma\gamma WW$ coupling, and no other vertices are involved in the unitary gauge at the lowest order [3]. The sensitivity to the γWW coupling is comparable and complementary to the reactions $e^+e^- \rightarrow W^+W^-$ and $\gamma e^- \rightarrow W^-\nu$: the first process involves a mixture of the γWW and the ZWW coupling, the second – involves the γWW alone but is not such sensitive. Because the sensitivity to the $\gamma\gamma WW$ coupling is much larger in the considering process than the one in e^+e^- collisions, $\gamma\gamma \rightarrow W^+W^-$ is the ideal process to study this coupling.




Fig. 1. The Feynman diagrams for W^+W^- -production

The measurement of cross section and asymmetries of the process $\gamma\gamma \rightarrow W^+W^-Z$ is complimentary to the analysis of the production of vector-boson pairs.

We have the unique possibility to examine production of two and three vector bosons in $\gamma\gamma$ collisions through the reaction using beams of polarized and unpolarized photons.

- This process involves only interactions between the gauge bosons making more evident any deviation from predictions of the gauge structure by Standard Model.
- There is no tree-level contribution involving the Higgs boson with any uncertainties coming from the scalar sector.

We started from explicit expression for the amplitude of process $\gamma\gamma \rightarrow W^+W^-$

$$M = G_v \epsilon_\mu(k_1) \epsilon_\nu(k_2) \epsilon_\alpha(p_+) \epsilon_\beta(p_-) M_T^{\mu\nu\alpha\beta}, \quad (1)$$

where

$$M_T^{\mu\nu\alpha\beta} = \sum_{i=1}^3 M_i^{\mu\nu\alpha\beta}, \quad (2)$$

k_1, k_2, p_+, p_- are four-momenta of the γ, γ, W^+, W^- , respectively and $\epsilon_\mu(k_1), \epsilon_\nu(k_2), \epsilon_\alpha(p_+), \epsilon_\beta(p_-)$ – their polarizations.

$$G_v = e^3 \cot \theta_W.$$

It is convenient to define the triple-gauge-boson coupling in the following way:

$$\Gamma_3^{\alpha\beta\gamma}(P_1, P_2) = [(2P_1 + P_2)^\beta g^{\alpha\gamma} - (2P_2 + P_1)^\alpha g^{\beta\gamma} + (P_2 - P_1)^\gamma g^{\alpha\beta}], \quad (3)$$

the quartic-gauge-boson coupling as

$$\Gamma_4^{\mu\nu\alpha\beta} = g^{\mu\alpha} g^{\nu\beta} + g^{\mu\beta} g^{\nu\alpha} - 2g^{\mu\nu} g^{\alpha\beta}, \quad (4)$$

and the propagator tensor for the vector boson as

$$D^{\mu\nu}(k) = \frac{(g^{\mu\nu} - k^\mu k^\nu / m^2)}{k^2 - m^2}. \quad (5)$$

Using the expression defined above, the contributions of the three Feynman diagrams (see fig. 1) for the WW -production can be written as

$$M_1^{\mu\nu\alpha\beta} = \Gamma_3^{\mu\alpha\xi}(-k_1, p_+) D_{\xi\lambda}(p_+ - k_1) \Gamma_3^{\nu\beta\lambda}(-k_2, p_-), \quad (6)$$

$$M_2^{\mu\nu\alpha\beta} = \Gamma_3^{\mu\beta\xi}(-k_1, p_-) D_{\xi\lambda}(p_- - k_1) \Gamma_3^{\nu\alpha\lambda}(-k_2, p_+), \quad (7)$$

$$M_3^{\mu\nu\alpha\beta} = \Gamma_4^{\mu\nu\alpha\beta}. \quad (8)$$

Then we calculate cross sections for polarized beams and for all combinations of polarizations of final particles. The exact covariant expressions for all amplitudes of different Feynman diagrams (see fig. 1) are presented in the appendix (21) – (24).

3. Cross sections and asymmetries

We analyze the total cross sections of the considered process as well as different distributions of the final state vector bosons. We have evaluated the total cross section for the processes $\gamma\gamma \rightarrow W^+W^-$

$$\sigma = \int |M|^2 d\Gamma, \quad (9)$$

where $|M|$ can be defined through formula (21) – (24), the phase-space volume element is assumed as

$$d\Gamma = \frac{d^3p_+}{(2\pi)^3 2p_+^0} \frac{d^3p_-}{(2\pi)^3 2p_-^0} (2\pi)^4 \delta(k_1 + k_2 - p_+ - p_-).$$

The total cross sections of $\gamma\gamma \rightarrow W^+W^-$ at $\sqrt{s} \simeq 1$ TeV is about $\sigma = 3.89 \cdot 10^{-8} \text{ GeV}^{-2}$.

In order to reach the best understanding of these reactions we present here various differential cross-section as well as polarized asymmetries of two- and three-boson production:

$$\begin{aligned} A_1 &= \frac{d\sigma_{+,+,+,+}/dy_- - d\sigma_{+,-,+,+}/dy_-}{d\sigma_{+,+,+,+}/dy_- + d\sigma_{+,-,+,+}/dy_-}, \\ A_2 &= \frac{d\sigma_{+,+,+,-}/dy_- - d\sigma_{+,-,+, -}/dy_-}{d\sigma_{+,+,+,-}/dy_- + d\sigma_{+,-,+, -}/dy_-}, \\ A_3 &= \frac{d\sigma_{+,+,0,0}/dy_- - d\sigma_{+,-,0,0}/dy_-}{d\sigma_{+,+,0,0}/dy_- + d\sigma_{+,-,0,0}/dy_-}, \\ A_4 &= \frac{d\sigma_{+,+,unpol,unpol}/dy_- - d\sigma_{+,-,unpol,unpol}/dy_-}{d\sigma_{+,+,unpol,unpol}/dy_- + d\sigma_{+,-,unpol,unpol}/dy_-}. \end{aligned} \quad (10)$$

Here the following notations are introduced: $+$, $-$ denote right and left circular polarizations of gammas and final bosons, 0 means a longitudinal polarization, *unpol* relates to the case of unpolarized particles.

One can see the y_- -distribution ($y_- = (k_1, p_-)/S$) and different kinds of polarized asymmetries in fig. 5 – fig. 16.

Using these formulas we have constructed the Monte-Carlo event's generator [4]. The universal kinematical block of this generator allows us to produce 2-particles, 3-particles and even more particles final state in identical way. We have used this block to calculation above mentioned processes and we are going to use it to another similar calculations some others processes in TESLA kinematics. It is realized as a fast-working FORTRAN module which provides required number of events (up to several billions in realistic time) with a defined initial energy and final masses.

The considered process is very important to study quartic-boson vertex (see fig.4). The contribution of four-boson vertex to total cross section is investigated specially. Obviously, processes of $\gamma\gamma$ -scattering are sensitive to the four-boson vertex construction. The experiments with polarized particles give wide set of additional data for investigation of electroweak gauge models.

4. The lowest-order correction to the $\gamma\gamma \rightarrow W^+W^-$ process

Obviously the correct comparison of results for these two processes and precision analysis of data of the future experiments are impossible without calculation of full set the first-order radiative corrections, presented in figs. 2, 3 (see, for example, refs. [8] – [13]).

As a rule, the one-loop diagrams involving Higgs boson have been calculated in order to study the possible investigation of the Higgs boson via $\gamma\gamma \rightarrow H^* \rightarrow W^+W^-$. See, for example, refs. [5] – [7]. Only the channels of longitudinal W -boson production are sensitive to the Higgs mechanism, but the channels with W transfer are insensitive. This insensitivity to the Higgs sector render $\gamma\gamma \rightarrow W^+W^-$ even more suitable for the investigation of the self couplings.

So, at $\sqrt{s} \sim 1000 \text{ GeV}$ the radiative correction is about a value of Born cross section. In this work we focus on the most important QED one-loop corrections to $\gamma\gamma \rightarrow W^+W^-$.

The additional real photon contribution (see Feynman diagrams in fig. 2) and additional virtual photon contribution (see Feynman diagrams in fig. 3) are shown in figs. 18 – 21.

The radiative correction achieves maximal value in the kinematical region, where y_- is near 0.

There is a list of important expressions needed to calculate a radiative contribution to the processes in figs. 2 and 3. Here

$$|M_1 + M_2 + M_3 + M_4 + M_5 + M_6 + M_7|_{soft}^2 = \left(\frac{A_1^\gamma + A_2^\gamma}{2(p_+k)} + \frac{B_1^\gamma + B_2^\gamma}{2(p_-k)} \right)^2, \quad (11)$$

$$A_1^\gamma = C_{0\alpha\beta\mu\nu} [\Gamma_3^{\nu\beta\lambda}(k_2, -p_-, p_- - k_2) D_{\lambda\sigma}(p_- - k_2) \Gamma_3^{\mu\rho\sigma}(k_1, k_2 - p, -p_+) (\delta_{\rho\eta} - p_{+\rho} p_{+\eta} / m_W^2) \Gamma_3^{\gamma\eta\alpha}(0, p_+, -p_+) + \Gamma_3^{\gamma\rho\alpha}(0, -p_+, -P_+) (\delta_{\rho\eta} - p_{+\rho} p_{+\eta} / m_W^2) \Gamma_4^{\mu\nu\beta\eta}]_{\vec{p}_+=0}, \quad (12)$$

$$A_2^\gamma = C_{0\alpha\beta\mu\nu} [\Gamma_3^{\mu\beta\lambda}(k_1, -p_-, p_- - k_1) D_{\lambda\sigma}(p_- - k_1) \Gamma_3^{\nu\rho\sigma}(k_2, k_1 - p, -p_+) (\delta_{\rho\eta} - p_{+\rho} p_{+\eta} / m_W^2) \Gamma_3^{\gamma\eta\alpha}(0, p_+, -p_+) + \Gamma_3^{\gamma\rho\alpha}(0, -p_+, -P_+) (\delta_{\rho\eta} - p_{+\rho} p_{+\eta} / m_W^2) \Gamma_4^{\mu\nu\beta\eta}]_{\vec{p}_+=0}, \quad (13)$$

$$B_1^\gamma = C_{0\alpha\beta\mu\nu} [\Gamma_3^{\beta\eta}(0, -p_-, p_-) (\delta_{\rho\eta} - p_{-\rho} p_{-\eta} / m_W^2) \Gamma_3^{\mu\rho\sigma}(k_1, -p, -p_+ + k_2) D_{\lambda\sigma}(p_+ - k_2) \times \Gamma_3^{\nu\lambda\alpha}(k_2, p_+ - k_2, -p_+) + \Gamma_3^{\beta\rho}(0, -p_-, P_-) (\delta_{\rho\sigma} - p_{-\rho} p_{-\sigma} / m_W^2) \Gamma_4^{\mu\nu\alpha\sigma}]_{\vec{p}_+=0}, \quad (14)$$

$$B_2^\gamma = C_{0\alpha\beta\mu\nu} [\Gamma_3^{\beta\eta}(0, -p_-, p_-) (\delta_{\rho\eta} - p_{-\rho} p_{-\eta} / m_W^2) \Gamma_3^{\nu\rho\sigma}(k_2, -p, -p_+ + k_1) D_{\lambda\sigma}(p_+ - k_1) \times \Gamma_3^{\mu\lambda\alpha}(k_1, p_+ - k_1, -p_+) + \Gamma_3^{\beta\rho}(0, -p_-, P_-) (\delta_{\rho\sigma} - p_{-\rho} p_{-\sigma} / m_W^2) \Gamma_4^{\mu\nu\alpha\sigma}]_{\vec{p}_+=0},$$




Fig. 2. The radiative corrections for $\gamma\gamma \rightarrow W^+W^-$ due to additional real photon

After that one can define the expressions of D_1, D_2, D_3 .

$$\begin{aligned}
 D_1 &= (|A_1^\gamma + A_2^\gamma|^2 p_0^2 + |B_1^\gamma + B_2^\gamma|^2 m_W^2 + 2\text{Re}(A_1^\gamma + A_2^\gamma(B_1^\gamma + B_2^\gamma)p_-^0 m))(8m_W^2 p_-^{02})^{-1}, \\
 D_2 &= 2(|A_1^\gamma + A_2^\gamma|^2 p_0^2 + 2\text{Re}(A_1^\gamma + A_2^\gamma(B_1^\gamma + B_2^\gamma)p_-^0 m))(8m_W^2 p_-^{02})^{-1}, \\
 D_3 &= (|A_1^\gamma + A_2^\gamma|^2 p_0^2)(8m_W^2 p_-^{02})^{-1}.
 \end{aligned} \tag{15}$$

Note, really it is impossible to obtain information with high precision from experimental data without the high-order effects consideration [13]. The radiative correction is defined as

$$\begin{aligned}
 \int (\sum_i M_i)^2 \frac{d^3 k}{(2\pi)^3 2k^0} &= \int (\sum_i M_i)^2 \frac{d^{n-1} k}{(2\pi)^{n-1} 2k} = \int \frac{2\pi^{1/2n-1} k^{n-2} \sin^{n-3} \theta dk d\theta}{\Gamma(1/2n-1)(2\pi)^{n-1} 2k} (\sum_i M_i)^2, \\
 R.C. &= \frac{2\pi^{1/2n-1}}{(2\pi)^{n-1} \Gamma(1/2n-1)} \int_0^\omega k^{n-5} dk \int_0^\pi \sin^{n-3} \theta \frac{[D_1 - v \cos \theta D_2 + v^2 \cos^2 \theta D_3]}{(1 - v \cos \theta)^2},
 \end{aligned} \tag{16}$$




Fig. 3. The radiative corrections for $\gamma\gamma \rightarrow W^+W^-$ due to additional virtual photon and Z-boson

where $v = |\vec{p}_-|/p_-^0$. The first two terms in Loran's series are

$$\begin{aligned}
 C_{-1} &= \frac{1}{16\pi^2} \left[-\frac{2D_1}{1-v^2} - \frac{2D_3}{1-v^2} - \frac{2D_2}{(1-v^2)^2} - \frac{1}{v} \log\left(\frac{1+v}{1-v}\right) [D_2 + 2D_3] + 2D_3 \right], \\
 C_0 &= \frac{1}{16\pi^2} [(\log \omega - \log 2\sqrt{2} + \frac{1}{2}C)(16\pi^2 C_{-1}) + \frac{1}{2}(\frac{4\log 2}{1-v^2} + \frac{2}{1-v^2} \log \frac{1-v}{1+v}) \times \\
 &\times (D_1 - D_2 + D_3) - \frac{1}{v}(\log \frac{1-v}{1+v}(\log(1 - \frac{1}{v}) + \log(1 + \frac{1}{v}))) + \Phi[\frac{1+v}{1-v}] - \Phi[\frac{1-v}{1+v}] \times \\
 &\times (D_2 - 2D_3 - 2D_3(\log 2 - 1))].
 \end{aligned} \tag{17}$$

There are some integrals needed for the calculation of a virtual γ -contributions. This part

of the correction was calculated using integration package of the *Mathematica*4.0.

$$\begin{aligned}
I_1 &= \int \frac{d^n k}{(2\pi)^n} \frac{1}{k^2((p' - k)^2 - m_W^2)((p + k)^2 - m_W^2)} = \\
&= \frac{-i}{16\pi^2} \frac{1}{2} \left(\frac{1}{n-4} + \frac{1}{2} C - \log 2\sqrt{\pi} + \log m \right) - \frac{1}{2} (1 - \log 2 - \log(1 + \frac{pp'}{m_W^2})) \frac{1}{m_W^2 + pp'}, \\
I_\alpha &= \int \frac{d^n k}{(2\pi)^n} \frac{k_\alpha}{k^2((p' - k)^2 - m_W^2)((p + k)^2 - m_W^2)} = \\
&= \frac{1}{q^2} \frac{1}{2p(p + p')/m_W^2 + 1} \left[\left(\frac{2p(p + p')}{m_W^2} + 2 \right) \log \left(\frac{2p(p + p')}{m_W^2} + 2 \right) - \left(\frac{2p(p + p')}{m_W^2} + 1 \right) \right] q_\alpha, \\
I_{\alpha\beta} &= \int \frac{d^n k}{(2\pi)^n} \frac{k_{\alpha\beta}}{k^2((p' - k)^2 - m_W^2)((p + k)^2 - m_W^2)} = \\
&= A\delta_{\alpha\beta} + B(p_\alpha p_\beta + p'_\alpha p'_\beta) - C(p_\alpha p'_\beta + p_\beta p'_\alpha),
\end{aligned} \tag{18}$$

where A, B, C are given by following equations

$$\begin{aligned}
A &= \frac{L^{(n)} + 2(pp')(B + C)}{4}, \\
B &= \frac{J_1 + 2(-m_W^2 - 2(pp'))C}{2(pp')}, \\
C &= \frac{2J_2 - L^{(n)} - J_1[-m_W^2/4 - 3/2(pp')]/(2(pp'))}{7/4(pp') + (-m_W^2 - 2(pp'))/(pp')(-m_W^2/4 + 3/2(pp'))},
\end{aligned} \tag{19}$$

$$\begin{aligned}
L^{(n)} &= \frac{-2i}{16\pi^2} \left[\left(\frac{1}{n-4} + \log m_W - \log 2\sqrt{\pi} + \frac{1}{2} C \right) + \frac{1}{\chi} ((\chi + 1) \log(\chi + 1) - \chi) \right], \\
J_1 &= D_0^{(n)} - D_1^{(n)} + \frac{p^2 - m_W^2}{q^2} \frac{1}{\chi} ((\chi + 1) \log(\chi + 1) - \chi), \\
J_2 &= D_2^{(n)} - \frac{p^2 - m_W^2}{q^2} \frac{1}{\chi} ((\chi + 1) \log(\chi + 1) - \chi), \\
D_0^{(n)} &= \frac{-i}{16\pi^2} \left[\left(\frac{1}{n-4} + \frac{1}{2} C - 1 \right) + \log m_W \right], \\
D_1^{(n)} &= \frac{-i}{16\pi^2} \left[\left(\frac{1}{n-4} + \log m_W - \log 2\sqrt{\pi} + \frac{1}{2} C \right) + \frac{1}{\chi^2} \left(\frac{(\chi + 1)^2}{2} \log(\chi + 1) - \right. \right. \\
&\quad \left. \left. - \frac{(\chi + 1)^2}{4} + \frac{1}{4} - (\chi + 1) \log(\chi + 1) + \chi \right) \right], \\
D_2^{(n)} &= \frac{-2i}{16\pi^2} \left[\left(\frac{1}{n-4} + \log m_W - \log 2\sqrt{\pi} + \frac{1}{2} C \right) + \frac{1}{\chi} ((\chi + 1) \log(\chi + 1) - \chi) - \right. \\
&\quad \left. - \frac{1}{2} \left(\frac{1}{\chi^2} \left(\frac{(\chi + 1)^2}{2} \log(\chi + 1) - \frac{(\chi + 1)^2}{4} + \frac{1}{4} - (\chi + 1) \log(\chi + 1) + \chi \right) \right) \right], \\
\chi &= 2p(p + p')/m_W^2 + 1,
\end{aligned} \tag{20}$$

where C - Euler's constant.

Appendix A

Using expressions (3) – (8) for the Feynman diagrams figs. 1 – 3 we have obtained the following matrix element for $\gamma\gamma \rightarrow W^+W^-$:

$$M = G_v \sum_{i=1}^3 M_i, \quad (21)$$

where M_i ($i = 1, 3$) are matrix elements related to the Feynman diagrams according to i .

$$\begin{aligned}
M_1 = & \frac{1}{(k_1 - p_+)^2 - m^2} (p_- \cdot \epsilon_2 \epsilon_- \cdot \epsilon_1 p_+ \cdot \epsilon_+ - p_- \cdot \epsilon_1 \epsilon_- \cdot \epsilon_2 p_+ \cdot \epsilon_+ - 2 p_- \cdot \epsilon_2 \epsilon_- \cdot \epsilon_+ p_+ \cdot \epsilon_1 + \\
& 2 p_- \cdot \epsilon_+ \epsilon_- \cdot \epsilon_2 p_+ \cdot \epsilon_1 + \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot p_+ p_+ \cdot \epsilon_+ p_+ \cdot \epsilon_1}{m^2} - \\
& \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot k_1 p_+ \cdot \epsilon_+ p_+ \cdot \epsilon_1}{m^2} - \frac{p_- \cdot p_+ \epsilon_- \cdot \epsilon_2 p_+ \cdot \epsilon_+ p_+ \cdot \epsilon_1}{m^2} + \\
& \frac{p_- \cdot k_1 \epsilon_- \cdot \epsilon_2 p_+ \cdot \epsilon_+ p_+ \cdot \epsilon_1}{m^2} - \frac{\epsilon_- \cdot \epsilon_2 p_+ \cdot \epsilon_+ p_+ \cdot k_2 p_+ \cdot \epsilon_1}{m^2} - \\
& \epsilon_- \cdot \epsilon_1 p_+ \cdot \epsilon_+ p_+ \cdot \epsilon_2 + 2 \epsilon_- \cdot \epsilon_+ p_+ \cdot \epsilon_1 p_+ \cdot \epsilon_2 + \frac{\epsilon_- \cdot k_2 p_+ \cdot \epsilon_+ p_+ \cdot \epsilon_1 p_+ \cdot \epsilon_2}{m^2} - \\
& 2 p_- \cdot \epsilon_2 \epsilon_- \cdot \epsilon_1 \epsilon_+ \cdot k_1 + 2 p_- \cdot \epsilon_1 \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot k_1 + 2 \epsilon_- \cdot \epsilon_1 p_+ \cdot \epsilon_2 \epsilon_+ \cdot k_1 + \\
& 2 \epsilon_- \cdot \epsilon_2 p_+ \cdot \epsilon_1 \epsilon_+ \cdot k_2 + p_- \cdot \epsilon_2 \epsilon_- \cdot p_+ \epsilon_+ \cdot \epsilon_1 + p_- \cdot \epsilon_2 \epsilon_- \cdot k_1 \epsilon_+ \cdot \epsilon_1 - \\
& p_- \cdot p_+ \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 - p_- \cdot k_1 \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 - \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot p_+ p_+ \cdot p_+ \cdot \epsilon_+ \cdot \epsilon_1}{m^2} + \\
& \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot k_1 p_+ \cdot p_+ \cdot \epsilon_+ \cdot \epsilon_1}{m^2} + \frac{p_- \cdot p_+ \epsilon_- \cdot \epsilon_2 p_+ \cdot p_+ \cdot \epsilon_+ \cdot \epsilon_1}{m^2} - \\
& \frac{p_- \cdot k_1 \epsilon_- \cdot \epsilon_2 p_+ \cdot p_+ \cdot \epsilon_+ \cdot \epsilon_1}{m^2} - \epsilon_- \cdot \epsilon_2 p_+ \cdot k_2 \epsilon_+ \cdot \epsilon_1 + \\
& \frac{\epsilon_- \cdot \epsilon_2 p_+ \cdot p_+ \cdot k_2 \epsilon_+ \cdot \epsilon_1}{m^2} - 2 \epsilon_- \cdot k_1 p_+ \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 + \\
& \epsilon_- \cdot k_2 p_+ \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 - \frac{\epsilon_- \cdot k_2 p_+ \cdot p_+ \cdot p_+ \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1}{m^2} - \\
& 2 \epsilon_- \cdot p_+ p_+ \cdot \epsilon_1 \epsilon_+ \cdot \epsilon_2 + 2 \epsilon_- \cdot k_1 p_+ \cdot \epsilon_1 \epsilon_+ \cdot \epsilon_2 - 2 \epsilon_- \cdot k_2 p_+ \cdot \epsilon_1 \epsilon_+ \cdot \epsilon_2 + \\
& \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot p_+ \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_1}{m^2} - \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot k_1 \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_1}{m^2} - \\
& \frac{p_- \cdot p_+ \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_1}{m^2} + \frac{p_- \cdot k_1 \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_1}{m^2} - \\
& \frac{\epsilon_- \cdot \epsilon_2 p_+ \cdot k_2 \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_1}{m^2} + \frac{\epsilon_- \cdot k_2 p_+ \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_1}{m^2} + \\
& \frac{\epsilon_- \cdot \epsilon_2 p_+ \cdot \epsilon_+ p_+ \cdot \epsilon_1 k_1 \cdot k_2}{m^2} - \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_2 - \\
& \frac{\epsilon_- \cdot \epsilon_2 p_+ \cdot p_+ \cdot \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_2}{m^2} + \frac{\epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 k_1 \cdot k_1 k_1 \cdot k_2}{m^2} + \\
& p_- \cdot \epsilon_2 \epsilon_- \cdot \epsilon_+ k_1 \cdot \epsilon_1 - p_- \cdot \epsilon_+ \epsilon_- \cdot \epsilon_2 k_1 \cdot \epsilon_1 - \epsilon_- \cdot \epsilon_+ p_+ \cdot \epsilon_2 k_1 \cdot \epsilon_1 - \\
& \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot p_+ \epsilon_+ \cdot k_1 k_1 \cdot \epsilon_1}{m^2} + \frac{p_- \cdot \epsilon_2 \epsilon_- \cdot k_1 \epsilon_+ \cdot k_1 k_1 \cdot \epsilon_1}{m^2} + \\
& \frac{p_- \cdot p_+ \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot k_1 k_1 \cdot \epsilon_1}{m^2} - \frac{p_- \cdot k_1 \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot k_1 k_1 \cdot \epsilon_1}{m^2} + \\
& \frac{\epsilon_- \cdot \epsilon_2 p_+ \cdot k_2 \epsilon_+ \cdot k_1 k_1 \cdot \epsilon_1}{m^2} - \frac{\epsilon_- \cdot k_2 p_+ \cdot \epsilon_2 \epsilon_+ \cdot k_1 k_1 \cdot \epsilon_1}{m^2} - \\
& \epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot k_2 k_1 \cdot \epsilon_1 + \epsilon_- \cdot p_+ \epsilon_+ \cdot \epsilon_2 k_1 \cdot \epsilon_1 - \epsilon_- \cdot k_1 \epsilon_+ \cdot \epsilon_2 k_1 \cdot \epsilon_1 + \\
& \epsilon_- \cdot k_2 \epsilon_+ \cdot \epsilon_2 k_1 \cdot \epsilon_1 - \frac{\epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot k_1 k_1 \cdot k_2 k_1 \cdot \epsilon_1}{m^2} + \epsilon_- \cdot \epsilon_1 p_+ \cdot \epsilon_+ k_1 \cdot \epsilon_2 -
\end{aligned} \quad (22)$$

$$\begin{aligned}
& 2 \epsilon_- \epsilon_+ p_+ \epsilon_1 k_1 \epsilon_2 - \frac{\epsilon_- k_2 p_+ \epsilon_+ p_+ \epsilon_1 k_1 \epsilon_2}{m^2} - 2 \epsilon_- \epsilon_1 \epsilon_+ k_1 k_1 \epsilon_2 + \\
& 2 \epsilon_- p_+ \epsilon_+ \epsilon_1 k_1 \epsilon_2 + \epsilon_- k_2 \epsilon_+ \epsilon_1 k_1 \epsilon_2 + \frac{\epsilon_- k_2 p_+ p_+ \epsilon_+ \epsilon_1 k_1 \epsilon_2}{m^2} - \\
& \frac{\epsilon_- k_2 \epsilon_+ \epsilon_1 k_1 k_1 k_1 \epsilon_2}{m^2} + \epsilon_- \epsilon_+ k_1 \epsilon_1 k_1 \epsilon_2 + \\
& \frac{\epsilon_- k_2 \epsilon_+ k_1 k_1 \epsilon_1 k_1 \epsilon_2}{m^2} - \epsilon_- \epsilon_2 p_+ \epsilon_+ k_2 \epsilon_1 + 2 \epsilon_- \epsilon_2 \epsilon_+ k_1 k_2 \epsilon_1 + \\
& \epsilon_- p_+ p_+ \epsilon_+ \epsilon_1 \epsilon_2 - \epsilon_- k_1 p_+ \epsilon_+ \epsilon_1 \epsilon_2 + \epsilon_- k_2 p_+ \epsilon_+ \epsilon_1 \epsilon_2 - \\
& 2 \epsilon_- p_+ \epsilon_+ k_1 \epsilon_1 \epsilon_2 + 2 \epsilon_- k_1 \epsilon_+ k_1 \epsilon_1 \epsilon_2 - 2 \epsilon_- k_2 \epsilon_+ k_1 \epsilon_1 \epsilon_2),
\end{aligned}$$

$$\begin{aligned}
M_2 = & \frac{1}{(k_1 - p_-)^2 - m^2} (2 p_- \epsilon_1 p_- \epsilon_2 \epsilon_- \epsilon_+ - 2 p_- \epsilon_+ p_- \epsilon_1 \epsilon_- \epsilon_2 + \frac{p_- \epsilon_- p_- \epsilon_+ p_- \epsilon_1 p_+ \epsilon_2}{m^2} - \\
& 2 p_- \epsilon_1 \epsilon_- \epsilon_+ p_+ \epsilon_2 + p_- \epsilon_+ \epsilon_- \epsilon_1 p_+ \epsilon_2 - \frac{p_- p_- p_- \epsilon_+ \epsilon_- \epsilon_1 p_+ \epsilon_2}{m^2} - \\
& 2 p_- \epsilon_2 \epsilon_- \epsilon_1 \epsilon_+ k_1 + 2 p_- \epsilon_1 \epsilon_- \epsilon_2 \epsilon_+ k_1 - \frac{p_- \epsilon_- p_- \epsilon_1 p_+ \epsilon_2 \epsilon_+ k_1}{m^2} + \\
& \epsilon_- \epsilon_1 p_+ \epsilon_2 \epsilon_+ k_1 + \frac{p_- p_- \epsilon_- \epsilon_1 p_+ \epsilon_2 \epsilon_+ k_1}{m^2} + \frac{p_- \epsilon_- p_- \epsilon_1 p_- \epsilon_2 \epsilon_+ k_2}{m^2} + \\
& p_- \epsilon_2 \epsilon_- \epsilon_1 \epsilon_+ k_2 - \frac{p_- p_- p_- \epsilon_2 \epsilon_- \epsilon_1 \epsilon_+ k_2}{m^2} - 2 p_- \epsilon_1 \epsilon_- \epsilon_2 \epsilon_+ k_2 - \\
& p_- \epsilon_- p_- \epsilon_2 \epsilon_+ \epsilon_1 + 2 p_- \epsilon_2 \epsilon_- k_1 \epsilon_+ \epsilon_1 + p_- \epsilon_- p_+ \epsilon_2 \epsilon_+ \epsilon_1 - \\
& 2 \epsilon_- k_1 p_+ \epsilon_2 \epsilon_+ \epsilon_1 - \frac{p_- \epsilon_- p_- p_+ p_- \epsilon_1 \epsilon_+ \epsilon_2}{m^2} - \frac{p_- \epsilon_- p_- k_2 p_- \epsilon_1 \epsilon_+ \epsilon_2}{m^2} + \\
& \frac{2 p_- \epsilon_1 \epsilon_- p_+ \epsilon_+ \epsilon_2 + 2 p_- \epsilon_1 \epsilon_- k_2 \epsilon_+ \epsilon_2 - p_- p_+ \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2 +}{m^2} \\
& \frac{p_- p_- p_- p_+ \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2}{m^2} - p_- k_2 \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2 + \frac{p_- p_- p_- k_2 \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2}{m^2} + \\
& \frac{p_- \epsilon_- p_- \epsilon_1 p_+ k_1 \epsilon_+ \epsilon_2}{m^2} - \epsilon_- \epsilon_1 p_+ k_1 \epsilon_+ \epsilon_2 - \frac{p_- p_- \epsilon_- \epsilon_1 p_+ k_1 \epsilon_+ \epsilon_2}{m^2} - \\
& p_- \epsilon_- p_+ \epsilon_1 \epsilon_+ \epsilon_2 + 2 \epsilon_- k_1 p_+ \epsilon_1 \epsilon_+ \epsilon_2 + \frac{p_- \epsilon_+ \epsilon_- \epsilon_1 p_+ \epsilon_2 k_1 k_1}{m^2} - \\
& \frac{\epsilon_- \epsilon_1 p_+ \epsilon_2 \epsilon_+ k_1 k_1 k_1}{m^2} + \frac{p_- \epsilon_2 \epsilon_- \epsilon_1 \epsilon_+ k_2 k_1 k_1}{m^2} - \\
& \frac{p_- p_+ \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2 k_1 k_1}{m^2} - \frac{p_- k_2 \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2 k_1 k_1}{m^2} + \frac{\epsilon_- \epsilon_1 p_+ k_1 \epsilon_+ \epsilon_2 k_1 k_1}{m^2} + \\
& \frac{p_- \epsilon_- p_- \epsilon_1 \epsilon_+ \epsilon_2 k_1 k_2}{m^2} - \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2 k_1 k_2 - \frac{p_- p_- \epsilon_- \epsilon_1 \epsilon_+ \epsilon_2 k_1 k_2}{m^2} + \\
& \frac{\epsilon_- \epsilon_1 \epsilon_+ \epsilon_2 k_1 k_1 k_1 k_2}{m^2} - p_- \epsilon_2 \epsilon_- \epsilon_+ k_1 \epsilon_1 + p_- \epsilon_+ \epsilon_- \epsilon_2 k_1 \epsilon_1 + \\
& \epsilon_- \epsilon_+ p_+ \epsilon_2 k_1 \epsilon_1 - \frac{p_- \epsilon_+ \epsilon_- k_1 p_+ \epsilon_2 k_1 \epsilon_1}{m^2} - \epsilon_- \epsilon_2 \epsilon_+ k_1 k_1 \epsilon_1 + \\
& \frac{\epsilon_- k_1 p_+ \epsilon_2 \epsilon_+ k_1 k_1 \epsilon_1}{m^2} - \frac{p_- \epsilon_2 \epsilon_- k_1 \epsilon_+ k_2 k_1 \epsilon_1}{m^2} + \epsilon_- \epsilon_2 \epsilon_+ k_2 k_1 \epsilon_1 - \\
& \epsilon_- p_+ \epsilon_+ \epsilon_2 k_1 \epsilon_1 + \frac{p_- p_+ \epsilon_- k_1 \epsilon_+ \epsilon_2 k_1 \epsilon_1}{m^2} + \frac{p_- k_2 \epsilon_- k_1 \epsilon_+ \epsilon_2 k_1 \epsilon_1}{m^2} - \\
& \epsilon_- k_2 \epsilon_+ \epsilon_2 k_1 \epsilon_1 - \frac{\epsilon_- k_1 p_+ k_1 \epsilon_+ \epsilon_2 k_1 \epsilon_1}{m^2} - \frac{\epsilon_- k_1 \epsilon_+ \epsilon_2 k_1 k_2 k_1 \epsilon_1}{m^2} - \\
& 2 p_- \epsilon_1 \epsilon_- \epsilon_+ k_1 \epsilon_2 + 2 p_- \epsilon_+ \epsilon_- \epsilon_1 k_1 \epsilon_2 - \frac{p_- \epsilon_- p_- \epsilon_1 \epsilon_+ k_2 k_1 \epsilon_2}{m^2} + \\
& \epsilon_- \epsilon_1 \epsilon_+ k_2 k_1 \epsilon_2 + \frac{p_- p_- \epsilon_- \epsilon_1 \epsilon_+ k_2 k_1 \epsilon_2}{m^2} + p_- \epsilon_- \epsilon_+ \epsilon_1 k_1 \epsilon_2 - \\
& 2 \epsilon_- k_1 \epsilon_+ \epsilon_1 k_1 \epsilon_2 - \frac{\epsilon_- \epsilon_1 \epsilon_+ k_2 k_1 k_1 k_1 \epsilon_2}{m^2} + \epsilon_- \epsilon_+ k_1 \epsilon_1 k_1 \epsilon_2 +
\end{aligned} \tag{23}$$

$$\begin{aligned} & \frac{\epsilon_- \cdot k_1 \epsilon_+ \cdot k_2 k_1 \cdot \epsilon_1 k_1 \cdot \epsilon_2}{m^2} - p_- \cdot \epsilon_- \epsilon_+ \cdot \epsilon_2 k_2 \cdot \epsilon_1 + 2 \epsilon_- \cdot k_1 \epsilon_+ \cdot \epsilon_2 k_2 \cdot \epsilon_1 + \\ & p_- \cdot \epsilon_- p_- \cdot \epsilon_+ \epsilon_1 \cdot \epsilon_2 - 2 p_- \cdot \epsilon_+ \epsilon_- \cdot k_1 \epsilon_1 \cdot \epsilon_2 - p_- \cdot \epsilon_- \epsilon_+ \cdot k_1 \epsilon_1 \cdot \epsilon_2 + \\ & 2 \epsilon_- \cdot k_1 \epsilon_+ \cdot k_1 \epsilon_1 \cdot \epsilon_2 + p_- \cdot \epsilon_- \epsilon_+ \cdot k_2 \epsilon_1 \cdot \epsilon_2 - 2 \epsilon_- \cdot k_1 \epsilon_+ \cdot k_2 \epsilon_1 \cdot \epsilon_2), \end{aligned}$$

$$M_3 = -\epsilon_- \cdot \epsilon_2 \epsilon_+ \cdot \epsilon_1 - \epsilon_- \cdot \epsilon_1 \epsilon_+ \cdot \epsilon_2 + 2 \epsilon_- \cdot \epsilon_+ \epsilon_1 \cdot \epsilon_2 \quad (24)$$

k_1, k_2, p_+, p_- denote four-momenta of γ, γ, W^+, W^- , respectively, and $\epsilon_1, \epsilon_2, \epsilon_+, \epsilon_-$ – cooresponding polarizations.

Appendix B




Fig. 4. The ratio $d\sigma_{1++}^{++}/d\sigma_{2++}^{++}$, where $d\sigma_{1(2)++}^{++}$ – 4-boson vertex is included (isn't included)




Fig. 5. The differential cross section $d\sigma_{++++}/dy_-$




Fig. 6. The differential cross section $d\sigma_{+,-,+,+}/dy_-$




Fig. 7. The asymmetry A_1




Fig. 8. The differential cross section $d\sigma_{+,+,+,-}/dy_-$




Fig. 9. The differential cross section $d\sigma_{+,-,+, -}/dy_-$




Fig. 10. The asymmetry A_2




Fig. 11. The differential cross section $d\sigma_{+,+,0,0}/dy_-$




Fig. 12. The differential cross section $d\sigma_{+,-,0,0}/dy_-$




Fig. 13. The asymmetry A_3




Fig. 14. The differential cross section $d\sigma_{+,+,unpol}/dy_-$




Fig. 15. The differential cross section $d\sigma_{+,-,unpol}/dy_-$




Fig. 16. The asymmetry A_4 .





Fig. 17. The differential cross section in Fig. 18. The radiative correction for Born approximation $d\sigma_{+, -, +, +}/dy^-$. $d\sigma_{+, -, +, +}/dy^-$.





Fig. 19. The differential cross section in Fig. 20. The radiative correction for Born approximation $d\sigma_{+, +, +, -}/dy^-$. $d\sigma_{+, +, +, -}/dy^-$.

References

- [1] I. Marfin, V. Mossolov, T. Shishkina hep-ph/0304250




Fig. 21. The differential cross section in Fig. 22. The radiative correction for Born approximation $d\sigma_{+,+,0,0}/dy_-$. $d\sigma_{+,+,0,0}/dy_-$.

- [2] TESLA Technical Design Report Part 6. Chapter 1. Photon collider at TESLA, hep-ex/0108012.
- [3] F. Brandt et al., Nucl. Phys. B450 (1994) p.223. hep-ph/9308353.
- [4] S. Weinzierl, NIKHEF-00-012. hep-ph/0006269.
- [5] M.Shifman et al. Sov J. Nucl. Phys. 30(1979) p.711.
- [6] E.Boos et al. Phys. Lett. B275(1992) p.164.
- [7] H.Veltman, Z.Phys. C62(1994) p.235.
- [8] M. Roth, A. Denner Nucl. Phys. B479 (1996) p.495. hep-ph/9605420.
- [9] A. Denner, S. Pozzorini Eur. Phys. J. C18 (2001) p.461. hep-ph/0010201
- [10] A. Denner, S. Pozzorini Eur. Phys. J. C21 (2001) p.63. hep-ph/0104127.
- [11] M. Melles Phys. Rept. 375 (2003) p.219. hep-ph/0104232.
- [12] F. Jegerlehner, M. Kalmykov, O. Veretin Nucl. Phys. B641 (2002) p.285. hep-ph/0105304.
- [13] B. Kniehl Int. J. Mod. Phys. A10 (1995) p.443. hep-ph/9410330.